arXiv:1304.3897v1 [math.CA] 14 Apr 2013

GENERALIZATION OF DIFFERENT TYPE INTEGRAL INEQUALITIES FOR $s-$CONVEX FUNCTIONS VIA FRACTIONAL INTEGRALS

İMDAT İŞCAN

Abstract. In this paper, a general integral identity for twice differentiable functions is derived. By using of this identity, the author establish some new estimates on Hermite-Hadamard type and Simpson type inequalities for $s-$convex via Riemann Liouville fractional integral.

1. Introduction

Following inequalities are well known in the literature as Hermite-Hadamard inequality and Simpson inequality respectively:

Theorem 1. *Let $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ be a convex function defined on the interval I of real numbers and $a,b\in I$ with $a<b$. The following double inequality holds*

$$f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int\limits_a^b f(x)dx\leq\frac{f(a)+f(b)}{2}. \tag{1.1}$$

Theorem 2. *Let $f:[a,b]\rightarrow\mathbb{R}$ be a four times continuously differentiable mapping on (a,b) and $\left\|f^{(4)}\right\|_{\infty}=\sup\limits_{x\in(a,b)}\left|f^{(4)}(x)\right|<\infty$. Then the following inequality holds:*

$$\left|\frac{1}{3}\left[\frac{f(a)+f(b)}{2}+2f\left(\frac{a+b}{2}\right)\right]-\frac{1}{b-a}\int\limits_a^b f(x)dx\right|\leq\frac{1}{2880}\left\|f^{(4)}\right\|_{\infty}(b-a)^2.$$

In [4], Hudzik and Maligranda considered among others the class of functions which are $s-$convex in the second sense.

Definition 1. *A function $f:[0,\infty)\rightarrow\mathbb{R}$ is said to be $s-$convex in the second sense if*

$$f(\alpha x+\beta y)\leq\alpha^s f(x)+\beta^s f(y)$$

for all $x,y\in[0,\infty)$, $\alpha,\beta\geq0$ with $\alpha+\beta=1$ and for some fixed $s\in(0,1]$. This class of $s-$convex functions in the second sense is usually denoted by K_s^2.

It can be easily seen that for $s=1$, $s-$convexity reduces to ordinary convexity of functions defined on $[0,\infty)$.

We give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

2000 *Mathematics Subject Classification.* 26A51, 26A33, 26D10.

Key words and phrases. Hermite–Hadamard inequality, Riemann–Liouville fractional integral, Simpson type inequalities, $s-$convex function.

Definition 2. *Let* $f\in L\left[a,b\right]$. *The Riemann-Liouville integrals* $J_{a^{+}}^{\alpha}f$ *and* $J_{b^{-}}^{\alpha}f$ *of oder* $\alpha>0$ *with* $a\geq 0$ *are defined by*

$$J_{a^{+}}^{\alpha}f(x)=\frac{1}{\Gamma(\alpha)}\int\limits_{a}^{x}\left(x-t\right)^{\alpha-1}f(t)dt,\ x>a$$

and

$$J_{b^{-}}^{\alpha}f(x)=\frac{1}{\Gamma(\alpha)}\int\limits_{x}^{b}\left(t-x\right)^{\alpha-1}f(t)dt,\ x<b$$

respectively, where $\Gamma(\alpha)$ *is the Gamma function defined by* $\Gamma(\alpha)=\int\limits_{0}^{\infty}e^{-t}t^{\alpha-1}dt$ *and* $J_{a^{+}}^{0}f(x)=J_{b^{-}}^{0}f(x)=f(x)$.

In the case of $\alpha=1$, the fractional integral reduces to the classical integral. Properties concerning this operator can be found [3, 5, 9].

In [11], Sarikaya et.al established the following theorem:

Theorem 3. *Let* $f:I\subset\mathbb{R}\rightarrow\mathbb{R}$ *be a twice differentiable mapping on* I° *such that* $f''\in L[a,b]$, *where* $a,b\in I$ *with* $a<b$. *If* $|f''|^{q}$ *is convex mapping on* $[a,b]$ *for* $q\geq 1$, *then the following inequality holds:*

$$\begin{aligned}&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{(b-a)}\int\limits_{a}^{b}f(x)dx\right|\\ &\leq \frac{\left(b-a\right)^{2}}{162}\left\{\left(\frac{59\left|f''\left(a\right)\right|^{q}+133\left|f''\left(b\right)\right|^{q}}{192}\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\frac{59\left|f''\left(b\right)\right|^{q}+133\left|f''\left(a\right)\right|^{q}}{192}\right)^{\frac{1}{q}}\right\}. \qquad (1.2)\end{aligned}$$

In [12], Bo-Yan Xi and Feng Qi establish some new integral inequalities of Hermite-Hadamard type for h-convex functions as follow:

Theorem 4. *Let* $I,J\subset\mathbb{R}$ *be intervals,* $(0,1)\subset J$, *and* $h:J\rightarrow\left[0,\infty\right)$. *Let* $f:I\subset\mathbb{R}\rightarrow\mathbb{R}$ *be a twice differentiable mapping on* I° *such that* $f''\in L[a,b]$ *for* $a,b\in I$ *with* $a<b$. *If* $0\leq\lambda\leq 1$ *and* $|f''|^{q}$ *is an* $h-$*convex mapping on* $[a,b]$ *for* $q\geq 1$, *then*

$$\begin{aligned}&\left|\left(1-\lambda\right)f\left(\frac{a+b}{2}\right)+\lambda\left(\frac{f(a)+f(b)}{2}\right)-\frac{1}{(b-a)}\int\limits_{a}^{b}f(x)dx\right|\\ &\leq \frac{\left(b-a\right)^{2}}{16}\left[H_{1}(\lambda)\right]^{1-\frac{1}{q}}\left\{\left(\left|f''\left(\frac{a+b}{2}\right)\right|^{q}\int\limits_{0}^{1}t\left|2\lambda-t\right|h(t)dt+\left|f''\left(a\right)\right|^{q}\int\limits_{0}^{1}t\left|2\lambda-t\right|h(1-t)dt\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\left|f''\left(\frac{a+b}{2}\right)\right|^{q}\int\limits_{0}^{1}t\left|2\lambda-t\right|h(t)dt+\left|f''\left(b\right)\right|^{q}C\int\limits_{0}^{1}t\left|2\lambda-t\right|h(1-t)dt\right)^{\frac{1}{q}}\right\},\end{aligned}$$

where

$$H_1(\lambda)=\begin{cases}\frac{8\lambda^3-3\lambda+1}{3}, & 0\leq\lambda\leq\frac{1}{2}\\ \frac{3\lambda-1}{3}, & \frac{1}{2}<\lambda\leq 1\end{cases}$$

Let us consider the following special functions:
(1) The Beta function:

$$\beta\left(x,y\right)=\frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}=\int\limits_0^1 t^{x-1}\left(1-t\right)^{y-1}dt,\quad x,y>0,$$

(2) The incomplete Beta function:

$$\beta\left(a,x,y\right)=\int\limits_0^a t^{x-1}\left(1-t\right)^{y-1}dt,\quad 0<a<1,\ x,y>0,$$

(3) The hypergeometric function:

$${}_2F_1\left(a,b;c;z\right)=\frac{1}{\beta\left(b,c-b\right)}\int\limits_0^1 t^{b-1}\left(1-t\right)^{c-b-1}\left(1-zt\right)^{-a}dt,\ c>b>0,\ |z|<1\text{ (see [1]).}$$

In Theorem 4, if we take $h(t)=t^s, s\in(0,1]$, we have the following inequality

$$\begin{aligned}&\left|(1-\lambda)f\left(\frac{a+b}{2}\right)+\lambda\left(\frac{f(a)+f(b)}{2}\right)-\frac{1}{(b-a)}\int\limits_a^b f(x)dx\right|\\ \leq\ & \frac{(b-a)^2}{16}\left[H_1(\lambda)\right]^{1-\frac{1}{q}}\left\{\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q H_2(\lambda,s)+\left|f''\left(a\right)\right|^q H_3(\lambda,s)\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q H_2(\lambda,s)+\left|f''\left(b\right)\right|^q H_3(\lambda,s)\right)^{\frac{1}{q}}\right\},\end{aligned}\tag{1.3}$$

where

$$H_2(\lambda,s)=\begin{cases}\frac{2(2\lambda)^{s+3}}{(s+2)(s+3)}-\frac{2\lambda}{s+2}+\frac{1}{s+3}, & 0\leq\lambda\leq\frac{1}{2}\\ \frac{2\lambda}{s+2}-\frac{1}{s+3}, & \frac{1}{2}<\lambda\leq 1\end{cases},$$

and

$$H_3(\lambda,s)=\begin{cases}\left[\begin{array}{c}2\lambda\beta\left(2,s+1\right)-\beta\left(3,s+1\right)\\ +4\lambda\beta\left(2\lambda;2,s+1\right)-2\beta\left(2\lambda;3,s+1\right)\end{array}\right], & 0\leq\lambda\leq\frac{1}{2}\\ 2\lambda\beta\left(2,s+1\right)-\beta\left(3,s+1\right), & \frac{1}{2}<\lambda\leq 1\end{cases},$$

In [7], Park establihed some new inequalities of the Simpson's like and the Hermite-Hadamar-like type for $s-$convex functions as follows:

Theorem 5. *Let $f:I\subset[0,\infty)\to\mathbb{R}$ be a twice differentiable function on I° such that f'' is integrable on $[a,b]$, where $a,b\in I^\circ$ with $a<b$. If $|f''|^q$ is $s-$convex on $[a,b]$ for some fixed $s\in(0,1]$ and $q>1$, then for $r\geq 2$ the following inequality holds:*

(a)

$$\begin{aligned}(1.4)\qquad & \frac{16}{(b-a)^2}\left|H_a^b(f)(r)\right| \\ \leq\ & \left\{\left[\frac{r-2}{r}\right]^{2p+1}\beta\left(1+p,1+p\right)+\frac{2^{p+1}}{r^{p+1}\left(p+1\right)}\cdot{}_2F_1\left(-p,1;p+2;\frac{2}{r}\right)\right\}^{\frac{1}{p}} \\ & \times\left\{\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+\left|f''\left(a\right)\right|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+\left|f''\left(b\right)\right|^q}{s+1}\right)^{\frac{1}{q}}\right\},\end{aligned}$$

where

$$\begin{aligned}& H_a^b(f)(r) \\ =\ & \left(\frac{1}{2}-\frac{1}{r}\right)\left\{\frac{f(a)+f(b)}{2}\right\}+\left(\frac{1}{2}+\frac{1}{r}\right)f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx.\end{aligned}$$

(b)

$$\begin{aligned}& \frac{16}{(b-a)^2}\left|R_a^b(f)(r)\right| \\ \leq\ & \left(\frac{2}{r}\right)\cdot\left(\frac{{}_2F_1\left(-p,1;p+2;\frac{-r}{2}\right)}{(p+1)}\right)^{\frac{1}{p}} \\ & \times\left\{\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+\left|f''\left(a\right)\right|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+\left|f''\left(b\right)\right|^q}{s+1}\right)^{\frac{1}{q}}\right\},\end{aligned}$$

where

$$\begin{aligned}& R_a^b(f)(r) \\ =\ & \left(\frac{1}{2}+\frac{1}{r}\right)\left\{\frac{f(a)+f(b)}{2}\right\}+\left(\frac{1}{2}-\frac{1}{r}\right)f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx.\end{aligned}$$

In recent years, many athors have studied errors estimations for Hermite-Hadamard's inequality and Simpson's on the class of $s-$convex mappings in the second sense and $(\alpha,m)-$convex mappings; for refinements, counterparts, generalizations and new Simpson's type inequalities, see [2, 6, 7, 8, 10, 11, 12].

The main aim of this article is to establish new generalization of Hermite Hadamard-type and Simpson-type inequalities for functions whose absolute values of second derivatives are $s-$convex. To begin with, the author will derive a general integral identity for twice differentiable mappings. By using this integral equality, the author establish some new inequalities of the Simpson-like and the Hermite-Hadamard-like type for these functions.

2. Main Results

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable function on I°, the interior of I, throughout this section we will take

$$\begin{aligned}
& I_f\left(x,\lambda,\alpha,a,b\right) \\
= \; & (1-\lambda)\left[\frac{\left(x-a\right)^{\alpha}+\left(b-x\right)^{\alpha}}{b-a}\right]f(x)+\lambda\left[\frac{\left(x-a\right)^{\alpha}f(a)+\left(b-x\right)^{\alpha}f(b)}{b-a}\right] \\
& +\left(\frac{1}{\alpha+1}-\lambda\right)\left[\frac{\left(b-x\right)^{\alpha+1}-\left(x-a\right)^{\alpha+1}}{b-a}\right]f'(x)-\frac{\Gamma\left(\alpha+1\right)}{b-a}\left[J_{x^-}^{\alpha}f(a)+J_{x^+}^{\alpha}f(b)\right]
\end{aligned}$$

where $a,b \in I$ with $a<b$, $x \in [a,b]$, $\lambda \in [0,1]$, $\alpha>0$ and Γ is Euler Gamma function. In order to prove our main results we need the following identity.

Lemma 1. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a twice differentiable function on I° such that $f'' \in L[a,b]$, where $a,b \in I$ with $a<b$. Then for all $x \in [a,b]$, $\lambda \in [0,1]$ and $\alpha>0$ we have:*

$$\begin{aligned}
I_f\left(x,\lambda,\alpha,a,b\right) &= \frac{\left(x-a\right)^{\alpha+2}}{\left(\alpha+1\right)\left(b-a\right)}\int_0^1 t\left(\left(\alpha+1\right)\lambda-t^{\alpha}\right)f''\left(tx+\left(1-t\right)a\right)dt \\
&+\frac{\left(b-x\right)^{\alpha+2}}{\left(\alpha+1\right)\left(b-a\right)}\int_0^1 t\left(\left(\alpha+1\right)\lambda-t^{\alpha}\right)f''\left(tx+\left(1-t\right)b\right)dt.
\end{aligned} \tag{2.1}$$

Proof. By integration by parts twice and changing the variable, for $x \neq a$, we can state

$$\begin{aligned}
& \int_0^1 t\left(\left(\alpha+1\right)\lambda-t^{\alpha}\right)f''\left(tx+\left(1-t\right)a\right)dt \\
= \; & t\left(\left(\alpha+1\right)\lambda-t^{\alpha}\right)\left.\frac{f'\left(tx+\left(1-t\right)a\right)}{x-a}\right|_0^1-\frac{\alpha+1}{x-a}\int_0^1\left(\lambda-t^{\alpha}\right)f'\left(tx+\left(1-t\right)a\right)dt \\
= \; & \left(\left(\alpha+1\right)\lambda-1\right)\frac{f'\left(x\right)}{x-a}-\frac{\alpha+1}{x-a}\left[\left(\lambda-t^{\alpha}\right)\left.\frac{f\left(tx+\left(1-t\right)a\right)}{x-a}\right|_0^1+\int_0^1\alpha t^{\alpha-1}\frac{f\left(tx+\left(1-t\right)a\right)}{x-a}dt\right] \\
= \; & \left(\left(\alpha+1\right)\lambda-1\right)\frac{f'\left(x\right)}{x-a}+\frac{\alpha+1}{\left(x-a\right)^2}\left[\left(1-\lambda\right)f(x)+\lambda f(a)-\frac{\alpha}{\left(x-a\right)^{\alpha}}\int_a^x\left(u-a\right)^{\alpha-1}f(u)du\right] \\
= \; & \left(\left(\alpha+1\right)\lambda-1\right)\frac{f'\left(x\right)}{x-a}+\frac{\alpha+1}{\left(x-a\right)^2}\left[\left(1-\lambda\right)f(x)+\lambda f(a)-\frac{\Gamma\left(\alpha+1\right)}{\left(x-a\right)^{\alpha}}J_{x^-}^{\alpha}f(a)\right].
\end{aligned} \tag{2.2}$$

Similarly, for $x \neq b$, we get

$$\begin{aligned}
& \int_0^1 t\left(\left(\alpha+1\right)\lambda-t^{\alpha}\right)f''\left(tx+\left(1-t\right)b\right)dt \\
= \; & -\left(\left(\alpha+1\right)\lambda-1\right)\frac{f'\left(x\right)}{b-x}+\frac{\alpha+1}{\left(b-x\right)^2}\left[\left(1-\lambda\right)f(x)+\lambda f(b)-\frac{\Gamma\left(\alpha+1\right)}{\left(b-x\right)^{\alpha}}J_{x^+}^{\alpha}f(b)\right]
\end{aligned} \tag{2.3}$$

.Multiplying both sides of (2.2) and (2.3) by $\frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)}$ and $\frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)}$, respectively, and adding the resulting identities, we obtain the desired result.

For $x=a$ and $x=b$,the identities

$$I_f\left(a,\lambda,\alpha,a,b\right)=\frac{(b-a)^{\alpha+1}}{(\alpha+1)}\int_0^1 t\left((\alpha+1)\lambda-t^{\alpha}\right)f''\left(ta+(1-t)b\right)dt,$$

and

$$I_f\left(b,\lambda,\alpha,a,b\right)=\frac{(b-a)^{\alpha+1}}{(\alpha+1)}\int_0^1 t\left((\alpha+1)\lambda-t^{\alpha}\right)f''\left(tb+(1-t)a\right)dt,$$

can be proved by performing an integration by parts twice in the integrals from the right side and changing the variable. □

Theorem 6. *Let $f:I\subset[0,\infty)\to\mathbb{R}$ be twice differentiable function on I° such that $f''\in L[a,b]$, where $a,b\in I^{\circ}$ with $a<b$. If $|f''|^q$ is $s-$convex on $[a,b]$ for some fixed $s\in(0,1]$ and $q\geq 1$, then for $x\in[a,b]$, $\lambda\in[0,1]$ and $\alpha>0$ the following inequality for fractional integrals holds*

$$\begin{aligned}
&\left|I_f\left(x,\lambda,\alpha,a,b\right)\right| \\
&\leq C_1^{1-\frac{1}{q}}\left(\alpha,\lambda\right)\left\{\frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)}\left(\left|f''(x)\right|^q C_2\left(\alpha,\lambda,s\right)+\left|f''(a)\right|^q C_3\left(\alpha,\lambda,s\right)\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)}\left(\left|f''(x)\right|^q C_2\left(\alpha,\lambda,s\right)+\left|f''(b)\right|^q C_3\left(\alpha,\lambda,s\right)\right)^{\frac{1}{q}}\right\},
\end{aligned}\tag{2.4}$$

where

$$\begin{aligned}
C_1\left(\alpha,\lambda\right) &= \begin{cases}\frac{\alpha[(\alpha+1)\lambda]^{1+\frac{2}{\alpha}}}{\alpha+2}-\frac{(\alpha+1)\lambda}{2}+\frac{1}{\alpha+2}, & 0\leq\lambda\leq\frac{1}{\alpha+1}\\ \frac{(\alpha+1)\lambda}{2}-\frac{1}{\alpha+2}, & \frac{1}{\alpha+1}<\lambda\leq 1\end{cases}, \\
C_2\left(\alpha,\lambda,s\right) &= \begin{cases}\frac{2\alpha[(\alpha+1)\lambda]^{\frac{\alpha+s+2}{\alpha}}}{(s+2)(\alpha+s+2)}-\frac{(\alpha+1)\lambda}{s+2}+\frac{1}{\alpha+s+2}, & 0\leq\lambda\leq\frac{1}{\alpha+1}\\ \frac{(\alpha+1)\lambda}{s+2}-\frac{1}{\alpha+s+2}, & \frac{1}{\alpha+1}<\lambda\leq 1\end{cases}, \\
C_3\left(\alpha,\lambda,s\right) &= \begin{cases}\begin{array}{c}\beta\left(\alpha+2,s+1\right)-(\alpha+1)\lambda\beta\left(2,s+1\right)\\ 2(\alpha+1)\lambda\beta\left(\left[(\alpha+1)\lambda\right]^{\frac{1}{\alpha}};2,s+1\right)-2\beta\left(\left[(\alpha+1)\lambda\right]^{\frac{1}{\alpha}};\alpha+2,s+1\right)\end{array}, & 0\leq\lambda\leq\frac{1}{\alpha+1}\\ (\alpha+1)\lambda\beta\left(2,s+1\right)-\beta\left(\alpha+2,s+1\right), & \frac{1}{\alpha+1}<\lambda\leq 1\end{cases}.
\end{aligned}$$

Proof. From Lemma 1, property of the modulus and using the power-mean inequality we have

$$
\begin{aligned}
\left|I_f\left(x,\lambda,\alpha,a,b\right)\right| &\leq \frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)}\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''\left(tx+(1-t)a\right)\right|dt \\
&+\frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)}\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''\left(tx+(1-t)b\right)\right|dt \\
&\leq \frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)}\left(\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|dt\right)^{1-\frac{1}{q}} \\
&\times\left(\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''\left(tx+(1-t)a\right)\right|^{q}dt\right)^{\frac{1}{q}} \\
&+\frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)}\left(\int_0^1 t\left|t\left(\alpha+1\right)\lambda-t^{\alpha}\right|dt\right)^{1-\frac{1}{q}} \\
&\times\left(\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''\left(tx+(1-t)b\right)\right|^{q}dt\right)^{\frac{1}{q}}.
\end{aligned}
\tag{2.5}
$$

Since $|f'|^q$ is $s-$convex on $[a,b]$ we get

$$
\begin{aligned}
\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''\left(tx+(1-t)a\right)\right|^{q}dt &\leq \int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left(t^{s}\left|f''(x)\right|^{q}+(1-t)^{s}\left|f''(a)\right|^{q}\right)dt \\
&= \left|f''(x)\right|^{q}C_2\left(\alpha,\lambda,s\right)+\left|f''(a)\right|^{q}C_3\left(\alpha,\lambda,s\right),
\end{aligned}
\tag{2.6}
$$

$$
\begin{aligned}
\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''\left(tx+(1-t)b\right)\right|^{q}dt &\leq \int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left(t^{s}\left|f''(x)\right|^{q}+(1-t)^{s}\left|f''(b)\right|^{q}\right)dt \\
&= \left|f''(x)\right|^{q}C_2\left(\alpha,\lambda,s\right)+\left|f''(b)\right|^{q}C_3\left(\alpha,\lambda,s\right),
\end{aligned}
\tag{2.7}
$$

where we use the fact that

$$
\begin{aligned}
& C_3(\alpha,\lambda,s) \\
= & \int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|(1-t)^s\,dt \\
= & \left\{\begin{array}{ll}
\begin{array}{c}
(\alpha+1)\lambda \displaystyle\int_0^{[(\alpha+1)\lambda]^{\frac{1}{\alpha}}} t(1-t)^s\,dt-\int_0^{[(\alpha+1)\lambda]^{\frac{1}{\alpha}}} t^{\alpha+1}(1-t)^s\,dt \\
-(\alpha+1)\lambda \displaystyle\int_{[(\alpha+1)\lambda]^{\frac{1}{\alpha}}}^1 t(1-t)^s\,dt+\int_{[(\alpha+1)\lambda]^{\frac{1}{\alpha}}}^1 t^{\alpha+1}(1-t)^s\,dt
\end{array}, & 0\le\lambda\le\frac{1}{\alpha+1} \\
(\alpha+1)\lambda \displaystyle\int_0^1 t(1-t)^s\,dt-\int_0^1 t^{\alpha+1}(1-t)^s\,dt, & \frac{1}{\alpha+1}<\lambda\le 1
\end{array}\right. \\
= & \left\{\begin{array}{ll}
\begin{array}{c}
\beta(\alpha+2,s+1)-(\alpha+1)\lambda\beta(2,s+1) \\
2(\alpha+1)\lambda\beta\left([(\alpha+1)\lambda]^{\frac{1}{\alpha}};2,s+1\right)-2\beta\left([(\alpha+1)\lambda]^{\frac{1}{\alpha}};\alpha+2,s+1\right)
\end{array}, & 0\le\lambda\le\frac{1}{\alpha+1} \\
(\alpha+1)\lambda\beta(2,s+1)-\beta(\alpha+2,s+1), & \frac{1}{\alpha+1}<\lambda\le 1
\end{array}\right.,
\end{aligned}
$$

$$
\begin{aligned}
& C_2(\alpha,\lambda,s) \\
= & \int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|t^s\,dt \\
= & \left\{\begin{array}{ll}
\frac{2\alpha[(\alpha+1)\lambda]^{\frac{\alpha+s+2}{\alpha}}}{(s+2)(\alpha+s+2)}-\frac{(\alpha+1)\lambda}{s+2}+\frac{1}{\alpha+s+2}, & 0\le\lambda\le\frac{1}{\alpha+1} \\
\frac{(\alpha+1)\lambda}{s+2}-\frac{1}{\alpha+s+2}, & \frac{1}{\alpha+1}<\lambda\le 1
\end{array}\right.
\end{aligned}
$$

and

$$
\begin{aligned}
& \int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|dt \\
= & \left\{\begin{array}{ll}
\frac{\alpha[(\alpha+1)\lambda]^{1+\frac{2}{\alpha}}}{\alpha+2}-\frac{(\alpha+1)\lambda}{2}+\frac{1}{\alpha+2}, & 0\le\lambda\le\frac{1}{\alpha+1} \\
\frac{(\alpha+1)\lambda}{2}-\frac{1}{\alpha+2}, & \frac{1}{\alpha+1}<\lambda\le 1
\end{array}\right.
\end{aligned}
\tag{2.8}
$$

Hence, If we use (2.6), (2.7) and (2.8) in (2.5), we obtain the desired result. This completes the proof. □

Corollary 1. *In Theorem 6,*

(a) If we choose $s=1$, *then we get:*

$$
\begin{aligned}
& \left|I_f(x,\lambda,\alpha,a,b)\right| \\
\le\ & C_1^{1-\frac{1}{q}}(\alpha,\lambda)\left\{\frac{(x-a)^{\alpha+1}}{b-a}\left(\left|f''(x)\right|^q C_2(\alpha,\lambda,1)+\left|f''(a)\right|^q C_3(\alpha,\lambda,1)\right)^{\frac{1}{q}}\right. \\
& \left.+\frac{(b-x)^{\alpha+1}}{b-a}\left(\left|f''(x)\right|^q C_2(\alpha,\lambda,1)+\left|f''(b)\right|^q C_3(\alpha,\lambda,1)\right)^{\frac{1}{q}}\right\}.
\end{aligned}
$$

(b) If we choose $q=1$, then we get:

$$\begin{aligned}\left|I_f(x,\lambda,\alpha,a,b)\right| &\leq \left\{\frac{(x-a)^{\alpha+1}}{b-a}\left(\left|f''(x)\right|C_2(\alpha,\lambda,s)+\left|f''(a)\right|C_3(\alpha,\lambda,s)\right)\right.\\ &\left.\quad+\frac{(b-x)^{\alpha+1}}{b-a}\left(\left|f''(x)\right|C_2(\alpha,\lambda,s)+\left|f''(b)\right|C_3(\alpha,\lambda,s)\right)\right\}.\end{aligned}$$

(c) If we choose $x=\frac{a+b}{2}$, then we get:

$$\begin{aligned}&\left|\frac{2^{\alpha-1}}{(b-a)^{\alpha-1}}I_f\left(\frac{a+b}{2},\lambda,\alpha,a,b\right)\right|\\ =&\left|(1-\lambda)f\left(\frac{a+b}{2}\right)+\lambda\left(\frac{f(a)+f(b)}{2}\right)-\frac{\Gamma(\alpha+1)2^{\alpha-1}}{(b-a)^{\alpha}}\left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-}f(a)+J^{\alpha}_{\left(\frac{a+b}{2}\right)^+}f(b)\right]\right|\\ \leq&\frac{(b-a)^2}{8(\alpha+1)}C_1^{1-\frac{1}{q}}(\alpha,\lambda)\left\{\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q C_2(\alpha,\lambda,s)+\left|f''(a)\right|^q C_3(\alpha,\lambda,s)\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q C_2(\alpha,\lambda,s)+\left|f''(b)\right|^q C_3(\alpha,\lambda,s)\right)^{\frac{1}{q}}\right\}.\end{aligned}\tag{2.9}$$

(d) If we choose $x=\frac{a+b}{2}$ and $\lambda=\frac{1}{3}$, then we get:

$$\begin{aligned}&\left|\frac{2^{\alpha-1}}{(b-a)^{\alpha-1}}I_f\left(\frac{a+b}{2},\frac{1}{3},\alpha,a,b\right)\right|\\ =&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{\Gamma(\alpha+1)2^{\alpha-1}}{(b-a)^{\alpha}}\left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-}f(a)+J^{\alpha}_{\left(\frac{a+b}{2}\right)^+}f(b)\right]\right|\\ \leq&\frac{(b-a)^2}{8(\alpha+1)}C_1^{1-\frac{1}{q}}\left(\alpha,\frac{1}{3}\right)\left\{\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q C_2\left(\alpha,\frac{1}{3},s\right)+\left|f''(a)\right|^q C_3\left(\alpha,\frac{1}{3},s\right)\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q C_2\left(\alpha,\frac{1}{3},s\right)+\left|f''(b)\right|^q C_3\left(\alpha,\frac{1}{3},s\right)\right)^{\frac{1}{q}}\right\}.\end{aligned}$$

(e) If we choose $x=\frac{a+b}{2}$, $\lambda=\frac{1}{3}$, and $\alpha=1$, then we get:

$$\begin{aligned}&\left|I_f\left(\frac{a+b}{2},\frac{1}{3},1,a,b\right)\right|\\ =&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{(b-a)}\int_a^b f(x)dx\right|\\ \leq&\frac{(b-a)^2}{162}\left(\frac{81}{8}\right)^{\frac{1}{q}}\left\{\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q C_2\left(1,\frac{1}{3},s\right)+\left|f''(a)\right|^q C_3\left(1,\frac{1}{3},s\right)\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\left|f''\left(\frac{a+b}{2}\right)\right|^q C_2\left(1,\frac{1}{3},s\right)+\left|f''(b)\right|^q C_3\left(1,\frac{1}{3},s\right)\right)^{\frac{1}{q}}\right\}.\end{aligned}$$

(f) If we choose $x=\frac{a+b}{2}$ *and* $\lambda=0$*, then we get:*

$$\begin{aligned}
&\left|\frac{2^{\alpha-1}}{(b-a)^{\alpha-1}}I_f\left(\frac{a+b}{2},0,\alpha,a,b\right)\right| \\
=&\left|f\left(\frac{a+b}{2}\right)-\frac{\Gamma(\alpha+1)2^{\alpha-1}}{(b-a)^{\alpha}}\left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-}f(a)+J^{\alpha}_{\left(\frac{a+b}{2}\right)^+}f(b)\right]\right| \\
\leq&\frac{(b-a)^2}{8(\alpha+1)}\left(\frac{1}{\alpha+2}\right)^{1-\frac{1}{q}}\left\{\left[\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q}{\alpha+s+2}+|f''(a)|^q\beta(\alpha+2,s+1)\right]^{\frac{1}{q}}\right. \\
&\left.+\left[\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q}{\alpha+s+2}+|f''(b)|^q\beta(\alpha+2,s+1)\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

(g) If we choose $x=\frac{a+b}{2}$*,* $\lambda=0$*, and* $\alpha=1$*, then we get:*

$$\begin{aligned}
&\left|I_f\left(\frac{a+b}{2},0,1,a,b\right)\right| \\
=&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{(b-a)}\int_a^b f(x)dx\right| \\
\leq&\frac{(b-a)^2}{16}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left\{\left[\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q}{s+3}+|f''(a)|^q\beta(3,s+1)\right]^{\frac{1}{q}}\right. \\
&\left.+\left[\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q}{s+3}+|f''(b)|^q\beta(3,s+1)\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

(h) If we choose $x=\frac{a+b}{2}$ *and* $\lambda=1$*, then we get:*

$$\begin{aligned}
&\left|\frac{2^{\alpha-1}}{(b-a)^{\alpha-1}}I_f\left(\frac{a+b}{2},1,\alpha,a,b\right)\right| \\
=&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)2^{\alpha-1}}{(b-a)^{\alpha}}\left[J^{\alpha}_{\left(\frac{a+b}{2}\right)^-}f(a)+J^{\alpha}_{\left(\frac{a+b}{2}\right)^+}f(b)\right]\right| \\
\leq&\frac{(b-a)^2}{8(\alpha+1)}\left(\frac{\alpha(\alpha+3)}{2(\alpha+2)}\right)^{1-\frac{1}{q}} \\
&\times\left\{\left[\frac{\alpha(\alpha+s+3)\left|f''\left(\frac{a+b}{2}\right)\right|^q}{(s+2)(\alpha+s+2)}+|f''(a)|^q((\alpha+1)\beta(2,s+1)-\beta(\alpha+2,s+1))\right]^{\frac{1}{q}}\right. \\
&\left.+\left[\frac{\alpha(\alpha+s+3)\left|f''\left(\frac{a+b}{2}\right)\right|^q}{(s+2)(\alpha+s+2)}+|f''(b)|^q((\alpha+1)\beta(2,s+1)-\beta(\alpha+2,s+1))\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

(i) If we choose $x = \frac{a+b}{2}, \lambda = 1$ and $\alpha = 1$, then we get:

$$
\begin{aligned}
& \left| I_f\left(\frac{a+b}{2}, 1, 1, a, b\right) \right| \\
= & \left| \frac{f(a)+f(b)}{2} - \frac{1}{(b-a)} \int_a^b f(x)dx \right| \\
\leq & \frac{(b-a)^2}{16} \left(\frac{2}{3}\right)^{1-\frac{1}{q}} \\
& \times \left\{ \left[\frac{(s+4)\left|f''\left(\frac{a+b}{2}\right)\right|^q}{(s+2)(s+3)} + \left|f''(a)\right|^q \left(2\beta(2, s+1) - \beta(3, s+1)\right) \right]^{\frac{1}{q}} \right. \\
& \left. + \left[\frac{(s+4)\left|f''\left(\frac{a+b}{2}\right)\right|^q}{(s+2)(s+3)} + \left|f''(b)\right|^q \left(2\beta(2, s+1) - \beta(3, s+1)\right) \right]^{\frac{1}{q}} \right\}.
\end{aligned}
$$

Remark 1. *In (c) of Corollary 1, if we choose $\alpha = 1$, then the inequality (2.9) reduces to the inequality (1.3).*

Remark 2. *In (e) of Corollary 1, if we choose $s = 1$, we have the following Simpson type inequality*

$$
\begin{aligned}
& \left| \frac{1}{6}\left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{(b-a)} \int_a^b f(x)dx \right| \\
\leq & \frac{(b-a)^2}{16} \left(\frac{8}{81}\right)^{1-\frac{1}{q}} \left\{ \left(\frac{59}{972} \left| f''\left(\frac{a+b}{2}\right) \right|^q + \frac{27}{972} \left|f''(a)\right|^q \right)^{\frac{1}{q}} \right. \\
& \left. + \left(\frac{59}{972} \left| f''\left(\frac{a+b}{2}\right) \right|^q + \frac{27}{972} \left|f''(b)\right|^q \right)^{\frac{1}{q}} \right\}.
\end{aligned}
$$

which is better than the inequality (1.2).

Theorem 7. *Let $f : I \subset [0, \infty) \to \mathbb{R}$ be twice differentiable function on I° such that $f'' \in L[a, b]$, where $a, b \in I^\circ$ with $a < b$. If $|f''|^q$ is $s-$convex on $[a, b]$ for some fixed $s \in (0, 1]$ and $q > 1$, then for $x \in [a, b]$, $\lambda \in [0, 1]$ and $\alpha > 0$ the following inequality for fractional integrals holds*

$$
\begin{aligned}
& \left| I_f(x, \lambda, \alpha, a, b) \right| \\
\leq & C_4^{\frac{1}{p}}(\alpha, \lambda, p) \left\{ \frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)} \left(\frac{\left|f''(x)\right|^q + \left|f''(a)\right|^q}{s+1} \right)^{\frac{1}{q}} \right. \\
& \left. + \frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)} \left(\frac{\left|f''(x)\right|^q + \left|f''(b)\right|^q}{s+1} \right)^{\frac{1}{q}} \right\},
\end{aligned}
\tag{2.10}
$$

where $p=\frac{q}{q-1}$ *and*

$$
\begin{aligned}
&C_4(\alpha,\lambda,p)\\
=&\left\{\begin{array}{cc}
\frac{1}{p(\alpha+1)+1}, & \lambda=0\\
\left[\begin{array}{c}\frac{[(\alpha+1)\lambda]^{\frac{1+(\alpha+1)p}{\alpha}}}{\alpha}\beta\left(\frac{1+p}{\alpha},1+p\right)\\
+\frac{[1-(\alpha+1)\lambda]^{p+1}}{\alpha(p+1)}\cdot{}_2F_1\left(1-\frac{1+p}{\alpha},1;p+2;1-(\alpha+1)\lambda\right)\end{array}\right], & 0<\lambda\le\frac{1}{\alpha+1}\\
\frac{[(\alpha+1)\lambda]^{\frac{p(\alpha+1)+1}{\alpha}}}{\alpha}\beta\left(\frac{1}{(\alpha+1)\lambda};\frac{1+p}{\alpha},1+p\right), & \frac{1}{\alpha+1}<\lambda\le1
\end{array}\right. .
\end{aligned}
$$

Proof. From Lemma 1, property of the modulus and using the Hölder inequality we have

$$
\begin{aligned}
&|I_f(x,\lambda,\alpha,a,b)|\le\frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)}\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''(tx+(1-t)a)\right|dt\\
&+\frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)}\int_0^1 t\left|(\alpha+1)\lambda-t^{\alpha}\right|\left|f''(tx+(1-t)b)\right|dt\\
\le\;&\frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)}\left(\int_0^1 t^p\left|(\alpha+1)\lambda-t^{\alpha}\right|^p dt\right)^{\frac{1}{p}}\times\left(\int_0^1\left|f''(tx+(1-t)a)\right|^q dt\right)^{\frac{1}{q}}\\
&+\frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)}\left(\int_0^1 t^p\left|(\alpha+1)\lambda-t^{\alpha}\right|^p dt\right)^{\frac{1}{p}}\times\left(\int_0^1\left|f''(tx+(1-t)b)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}
\tag{2.11}
$$

Since $|f'|^q$ is $s-$convex on $[a,b]$ we get

$$
\begin{aligned}
\int_0^1\left|f''(tx+(1-t)a)\right|^q dt &\le \int_0^1\left(t^s\left|f''(x)\right|^q+(1-t)^s\left|f''(a)\right|^q\right)dt\\
&=\frac{\left|f''(x)\right|^q+\left|f''(a)\right|^q}{s+1},
\end{aligned}
\tag{2.12}
$$

$$
\begin{aligned}
\int_0^1\left|f''(tx+(1-t)b)\right|^q dt &\le \int_0^1\left(t^s\left|f''(x)\right|^q+(1-t)^s\left|f''(b)\right|^q\right)dt\\
&=\frac{\left|f''(x)\right|^q+\left|f''(b)\right|^q}{s+1},
\end{aligned}
\tag{2.13}
$$

and

$$\int_0^1 t^p \left|(\alpha+1)\lambda - t^\alpha\right|^p dt \tag{2.14}$$

$$= \begin{cases} \displaystyle\int_0^1 t^{(\alpha+1)p} dt & \lambda = 0 \\ \displaystyle\int_0^{[(\alpha+1)\lambda]^{\frac{1}{\alpha}}} t^p \left[(\alpha+1)\lambda - t^\alpha\right]^p dt + \int_{[(\alpha+1)\lambda]^{\frac{1}{\alpha}}}^1 t^p \left[t^\alpha - (\alpha+1)\lambda\right]^p dt, & 0 < \lambda \leq \frac{1}{\alpha+1} \\ \displaystyle\int_0^1 t^p \left[(\alpha+1)\lambda - t^\alpha\right]^p dt, & \frac{1}{\alpha+1} < \lambda \leq 1 \end{cases}$$

$$= \begin{cases} \frac{1}{p(\alpha+1)+1}, & \lambda = 0 \\ \left[\begin{array}{c} \frac{[(\alpha+1)\lambda]^{\frac{1+(\alpha+1)p}{\alpha}}}{\alpha} \beta\left(\frac{1+p}{\alpha}, 1+p\right) \\ + \frac{[1-(\alpha+1)\lambda]^{p+1}}{\alpha(p+1)} \cdot {}_2F_1\left(1 - \frac{1+p}{\alpha}, 1; p+2; 1-(\alpha+1)\lambda\right) \end{array}\right], & 0 < \lambda \leq \frac{1}{\alpha+1} \\ \frac{[(\alpha+1)\lambda]^{\frac{1+(\alpha+1)p}{\alpha}}}{\alpha} \beta\left(\frac{1}{(\alpha+1)\lambda}; \frac{1+p}{\alpha}, 1+p\right), & \frac{1}{\alpha+1} < \lambda \leq 1 \end{cases}$$

Hence, If we use (2.12), (2.13) and (2.14) in (2.11), we obtain the desired result. This completes the proof. □

Corollary 2. *In Theorem 7*

(a) If we choose $s = 1$, ten we get:

$$\begin{aligned} & \left|I_f(x, \lambda, \alpha, a, b)\right| \\ \leq\ & C_4^{\frac{1}{p}}(\alpha, \lambda, p) \left\{ \frac{(x-a)^{\alpha+2}}{(\alpha+1)(b-a)} \left(\frac{\left|f''(x)\right|^q + \left|f''(a)\right|^q}{2} \right)^{\frac{1}{q}} \right. \\ & \left. + \frac{(b-x)^{\alpha+2}}{(\alpha+1)(b-a)} \left(\frac{\left|f''(x)\right|^q + \left|f''(b)\right|^q}{2} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

(b) If we choose $x = \frac{a+b}{2}$, then we get:

$$\begin{aligned} & \left|I_f(x, \lambda, \alpha, a, b)\right| \\ \leq\ & C_4^{\frac{1}{p}}(\alpha, \lambda, p) \frac{(b-a)^{\alpha+1}}{(\alpha+1)2^{\alpha+2}} \left\{ \left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q + \left|f''(a)\right|^q}{s+1} \right)^{\frac{1}{q}} \right. \\ & \left. + \left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q + \left|f''(b)\right|^q}{s+1} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

(c) If we choose $x=\frac{a+b}{2},\ \lambda=\frac{1}{3}$*,then we get:*

$$\begin{aligned}&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{\Gamma(\alpha+1)\,2^{\alpha-1}}{(b-a)^{\alpha}}\left[J_{\left(\frac{a+b}{2}\right)^{-}}^{\alpha}f(a)+J_{\left(\frac{a+b}{2}\right)^{+}}^{\alpha}f(b)\right]\right|\\ \leq\ & C_4^{\frac{1}{p}}\left(\alpha,\frac{1}{3},p\right)\frac{(b-a)^2}{8(\alpha+1)}\left\{\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(a)|^q}{s+1}\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\}.\end{aligned}$$

(d) If we choose $x=\frac{a+b}{2},\ \lambda=\frac{1}{3}$*, and* $\alpha=1$*,then we get:*

$$\begin{aligned}&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{(b-a)}\int_a^b f(x)dx\right|\\ \leq\ & \frac{(b-a)^2}{16}C_4^{\frac{1}{p}}\left(1,\frac{1}{3},p\right)\left\{\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(a)|^q}{s+1}\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\},\end{aligned}$$

where

$$C_4\left(1,\frac{1}{3},p\right)=\left(\frac{2}{3}\right)^{1+2p}\beta(1+p,1+p)+\left(\frac{1}{3}\right)^{1+p}.{}_2F_1\left(-p,1;p+2;\frac{1}{3}\right).$$

(e) If we choose $x=\frac{a+b}{2}$ *and* $\lambda=0$*,then we get:*

$$\begin{aligned}&\left|f\left(\frac{a+b}{2}\right)-\frac{\Gamma(\alpha+1)\,2^{\alpha-1}}{(b-a)^{\alpha}}\left[J_{\left(\frac{a+b}{2}\right)^{-}}^{\alpha}f(a)+J_{\left(\frac{a+b}{2}\right)^{+}}^{\alpha}f(b)\right]\right|\\ \leq\ & \frac{(b-a)^2}{16}\left(\frac{1}{p(\alpha+1)+1}\right)^{\frac{1}{p}}\left\{\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(a)|^q}{s+1}\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\}.\end{aligned}$$

(f) If we choose $x=\frac{a+b}{2}$ *and* $\lambda=1$*,then we get:*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)\,2^{\alpha-1}}{(b-a)^{\alpha}}\left[J_{\left(\frac{a+b}{2}\right)^{-}}^{\alpha}f(a)+J_{\left(\frac{a+b}{2}\right)^{+}}^{\alpha}f(b)\right]\right|\\ \leq\ & \frac{(b-a)^2}{16}C_4^{\frac{1}{p}}(\alpha,1,p)\left\{\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(a)|^q}{s+1}\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+|f''(b)|^q}{s+1}\right)^{\frac{1}{q}}\right\},\end{aligned}$$

where

$$C_4(\alpha,1,p)=\frac{(1+\alpha)^{\frac{p(\alpha+1)+1}{\alpha}}}{\alpha}\beta\left(\frac{1}{1+\alpha};\frac{1+p}{\alpha},1+p\right).$$

(g) If we choose $x=\frac{a+b}{2}$, $\lambda=1$ *and* $\alpha=1$,*then we get:*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{(b-a)}\int\limits_a^b f(x)dx\right|\\ \leq\;&\frac{(b-a)^2}{4}\left(2\beta\left(\frac{1}{2};1+p,1+p\right)\right)^{\frac{1}{p}}\left\{\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+\left|f''(a)\right|^q}{s+1}\right)^{\frac{1}{q}}\right.\\ &\left.+\left(\frac{\left|f''\left(\frac{a+b}{2}\right)\right|^q+\left|f''(b)\right|^q}{s+1}\right)^{\frac{1}{q}}\right\}.\end{aligned}$$

Remark 3. *In (b) of Corollary 2, if we take* $\lambda=\frac{1}{2}-\frac{1}{r}, r\geq 2$, *and* $\alpha=1$, *then the inequality (2.10) reduces to the inequality (1.4).*

References

[1] M. Abramowitz, I.A. Stegun, eds., Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, Dover, New York (1965).

[2] M.I.Bhatti, M.Iqbal and S.S.Dragomir, Some new fractional integral Hermite-Hadamard type inequalities, RGMIA Res. Rep. Coll., 16 (2013), Article 2.

[3] R. Gorenflo, F. Mainardi, Fractional calculus; integral and differential equations of fractional order, Springer Verlag, Wien (1997), 223-276.

[4] H. Hudzik and L. Maligranda, Some remarks on $s-$convex functions, Aequationes Math, 48 (1994), 100-111.

[5] S. Miller and B. Ross, An introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, USA (1993).

[6] M. E. Özdemir, M. Avci and H. Kavurmaci, Hermite–Hadamard-type inequalities via $(\alpha,m)-$convexity, Computers and Mathematics with Applications 61 (2011), 2614–2620.

[7] J. Park, Hermite and Simpson-like type inequalities for functions whose second derivatives in absolute values at certain powera are $s-$convex, International Journal of Pure and Applied Mathematics, Vol. 78, No. 5, (2012), 587-604.

[8] J. Park,On the Simpson-like type inequalities for twice differentiable $(\alpha,m)-$convex mapping, International Journal of Pure and Applied Mathematics, Vol. 78, No. 5, (2012), 617–634.

[9] I. Podlubni, Fractional Differential Equations, Academic Press, San Diego, 1999.

[10] M.Z. Sarikaya and N. Aktan, On the generalization of some integral inequalities and their applications, Mathematical and Computer Modelling 54 (2011), 2175–2182.

[11] M.Z. Sarikaya, E. Set and M.E. Ozdemir, On new inequalities of Simpson's type for convex functions, RGMIA Res. Rep. Coll., 13, No. 2 (2010), Article 2.

[12] B.-Y. Xi, F. Qi, Some inequalities of Hermite-Hadamard typr for $h-$convex functions, Advances in Inequalities and Applications, 2 (2013), No.1, 1-15.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES AND ARTS, GIRESUN UNIVERSITY, GIRESUN, TURKEY

E-mail address: `imdat.iscan@giresun.edu.tr`